

# Transcriptional regulation of methanogenic metabolism in archaea

Katie E Shalvarjian<sup>1</sup> and Dipti D Nayak



Methanogenesis is a widespread metabolism of evolutionary and environmental importance that is likely to have originated on early Earth. Microorganisms that perform methanogenesis, termed methanogens, belong exclusively to the domain Archaea. Despite maintaining eukaryotic transcription machinery and homologs of bacterial regulators, archaeal transcription and gene regulation appear to be distinct from either domain. While genes involved in methanogenic metabolism have been identified and characterized, their regulation in response to both extracellular and intracellular signals is less understood. Here, we review recent reports on transcriptional regulation of methanogenesis using two model methanogens, *Methanococcus maripaludis* and *Methanosarcina acetivorans*, and highlight directions for future research in this nascent field.

## Address

Department of Molecular and Cell Biology, University of California, Berkeley, CA, USA

Corresponding author: Nayak, Dipti D ([dnayak@berkeley.edu](mailto:dnayak@berkeley.edu))

<sup>1</sup> Present Address: Department of Plant and Microbial Biology, 111 Koshland Hall, University of California, Berkeley, CA 94720.

Current Opinion in Microbiology 2021, 60:8–15

This review comes from a themed issue on Cell regulation

Edited by Sean Crosson and Aretha Fiebig

<https://doi.org/10.1016/j.mib.2021.01.005>

1369-5274/© 2021 Elsevier Ltd. All rights reserved.

## Introduction

Methanogens are a polyphyletic group of archaea that couple growth and energy conservation to the generation of methane gas through a process called methanogenesis [1–3]. These microbes are typically found in anoxic environments such as wetlands and sewage treatment plants, as well as the digestive tract of diverse hosts like termites, cows, and humans [4]. As the predominant source of methane, a potent greenhouse gas, methanogenic archaea rank as a highly influential group of organisms in the context of global climate regulation [1,5–7]. By now, many of the enzymes and respiratory complexes involved in methanogenesis have been identified and characterized in intricate biochemical detail [1,3]. Yet, very little is known about how

these genes are regulated in response to biotic or abiotic signals. As the climate crisis exacerbates, the need to understand how methanogens regulate their metabolism in response to environmental cues is crucial to forecast how they will adapt to and impact the warming planet.


Since all known methanogens are members of the Archaea, the third domain of Life, their transcriptional machinery contains elements related to that of Bacteria and Eukarya, while remaining distinct from either (Figure 1) [8,9]. For instance, the archaeal RNA polymerase (RNAP) is structurally and evolutionarily related to PolIII in *Saccharomyces cerevisiae* (Figure 1c) [10]. Similarly, transcription initiation relies on homologs of eukaryotic TATA-binding proteins (TBPs) and transcription factor B (TFB) binding to the TATA box and B-recognition element in the core promoter sequence, respectively (Figure 1a) [11,12]. However, following coordination of RNAP with the promoter, elongation proceeds akin to Bacteria, with transcription of the nascent mRNA coupled directly to its translation (Figure 1b) [13]. This is particularly relevant within the context of polarity as archaeal genomes are generally organized in operons [8]. Although homologs of the canonical, bacterial two component system (TCS) and some eukaryotic transcription factors have been identified in methanogens, the fundamental mechanisms of signal transduction and gene regulation in archaea are distinct and largely uncharacterized *in vivo* [14,15,16\*]. However, with the advent of new genetic techniques, these questions are becoming more feasible to address within the context of the cell.

This review summarizes current knowledge and identifies future directions for research related to the regulation of methanogenesis in *Methanococcus maripaludis* and *Methanosarcina acetivorans*. These well-studied species have emerged as genetically tractable model organisms that represent two distinct energy conservation strategies in methanogens. *M. maripaludis* lacks cytochromes and is restricted to growth on H<sub>2</sub> + CO<sub>2</sub> or formate [17]; whereas *M. acetivorans* contains cytochromes and has a broader substrate range, which can include acetate and methylated compounds like methanol or methylamines [18]. Together, these two species provide an ideal genetic framework to identify signaling cascades and transcription factors that regulate methanogenesis *in vivo*.

## Methanogens without cytochromes

Methanogens belonging to the Orders *Methanococcales*, *Methanobacteriales*, *Methanomicrobiales*, and *Methanopyrales*

Figure 1




Transcription in Methanogenic Archaea. **(a)** Two transcription factors, TBP (TATA binding protein; in orange) and TFB (transcription factor B; in orange-red), that bind to the TATA-box and the B-recognition element (BRE) in the promoter region respectively are required to recruit the RNA polymerase (in light blue) for transcription initiation in methanogenic archaea. A third transcription factor, Transcription factor E (TFE; in yellow) is not essential but enhances the recruitment of RNA polymerase *in vitro*. The 5' untranslated region (UTR) is highlighted in blue. **(b)** Genes in methanogenic archaea are arranged in operons and transcription of a multi-gene operon (in green) produces polycistronic RNA. Assembly of the corresponding polypeptide is facilitated by a translating ribosome (light purple). **(c)** Structural overview of RNA polymerase (RNAP) derived from a bacterium (*Escherichia coli*; PDB ID: 4YG2), an archaeon (*Thermococcus kodakarensis* PDB ID: 4QIW) and a eukaryote (Polymerase II from *Saccharomyces cerevisiae* PDB ID: 1WCM). Orthologous subunits of RNAP are shown using the same color.

within the *Euryarchaeota*, lack cytochromes and differ in their energy metabolism from members of *Methanosarcinales*. The vast majority of these cytochrome-lacking methanogens are restricted to growth on CO<sub>2</sub> with H<sub>2</sub> or formate as the electron donor [1,4]. However, a few notable exceptions can use methylated compounds as a carbon source and short-

chain alcohols, like ethanol, as an electron donor as discussed in [2,19,20]. Although methanogens with and without cytochromes share a conserved seven-step pathway for the step-wise reduction of CO<sub>2</sub> to methane, the two differ significantly in their energy conservation strategies [1]. Methanogens without cytochromes, like *M. maripaludis*, also lack the

Figure 2



Condition Specific Regulation of Methanogenesis in *Methanococcus maripaludis*. Steps highlighted with colored arrows denote upregulation of the corresponding enzyme based on substrate availability. **(a)** Transcriptomic analyses of methanogenesis in H<sub>2</sub>-replete and H<sub>2</sub>-limited conditions has shown that genes catalyzing the fourth and the fifth step in the sequential reduction of CO<sub>2</sub> to methane are transcriptionally regulated as a function of H<sub>2</sub> availability. Additionally, the F<sub>420</sub>-reducing hydrogenase complex (Fru) and formate dehydrogenase (Fdh1) are upregulated upon H<sub>2</sub> limitation. **(b)** During growth on formate, three steps of the seven-step pathway for CO<sub>2</sub> reduction are regulated as a function of substrate concentration. In addition, the two formate dehydrogenase orthologs are differently expressed: Fdh1 is expressed in formate replete media whereas Fdh2 is only expressed when formate concentrations are growth limiting.

quinone-like compound methanophenazine and rely on flavin-based electron bifurcation (FBEB) for the regeneration of coenzyme M (CoM) and coenzyme B (CoB) from the oxidized heterodisulfide (CoM-CoB) produced during the final step of methanogenesis [1,21–23]. As a result, the Na<sup>+</sup>-dependent N<sup>5</sup>-methyl-tetrahydromethanopterin:CoM methyltransferase (Mtr) singularly generates the ion gradient for ATP synthesis by the membrane-bound ATPase [1]. Until recently, *M. maripaludis* was the only genetically tractable member of this group of methanogens [24,25]. While its nitrogen regulon has been characterized and reviewed in detail in [26,27], regulation of methanogenesis in *M. maripaludis* has only been inferred from genome-wide transcriptomic data [28]. Therefore, despite extensive evidence for the regulation of methanogenic enzymes (as discussed below) transcriptional regulators involved in this process are yet to be characterized.

### H<sub>2</sub> dependent gene regulation

Whole cell transcriptomic analyses of *M. maripaludis* in chemostat cultures where growth was limited by H<sub>2</sub>

revealed extensive regulation in core methanogenesis genes [28–30]. Significantly, expression of metabolic genes that use F<sub>420</sub>, an NAD(P) analog used by methanogens as a redox carrier, is linked to extracellular H<sub>2</sub> levels [29]. Two enzymes can carry out the fourth step of the methanogenesis pathway: an F<sub>420</sub>-dependent tetrahydromethanopterin dehydrogenase (Mtd) or a H<sub>2</sub>-dependent tetrahydromethanopterin dehydrogenase (Hmd) (Figure 2a). As H<sub>2</sub> levels drop, transcripts for Mtd become more abundant relative to Hmd, favoring the use of F<sub>420</sub> as an intermediate under low H<sub>2</sub> conditions (Figure 2a) [30,31<sup>••</sup>]. Similarly, transcripts for the F<sub>420</sub>-dependent methylene tetrahydromethanopterin reductase (Mer) and F<sub>420</sub>-dependent hydrogenase (Fru) are more abundant under H<sub>2</sub>-limiting conditions (Figure 2a) [29]. This upregulation of genes encoding F<sub>420</sub>-dependent enzymes under H<sub>2</sub>-limiting conditions might be beneficial to sustain metabolic flux for methanogenesis fueled by reduced F<sub>420</sub> rather than H<sub>2</sub> [17,23,31<sup>••</sup>]. Similar studies with other methanogens like *Methanocaldococcus janaschii* and *Methanothermobacter thermoautotrophicum* ΔH, have also revealed substantial transcriptional regulation of Mtd

versus Hmd—along with other methanogenic enzymes—as a function of H<sub>2</sub> availability [32,33]. Altogether, while it is clear that there is an extensive H<sub>2</sub> regulon in *M. maripaludis*, a H<sub>2</sub> sensor and its downstream signaling cascade remains elusive.

### Formate dependent gene regulation

In methanogens without cytochromes, formate dehydrogenase (encoded by *fdhAB*) catalyzes the oxidation of formate to CO<sub>2</sub> with the concomitant reduction of F<sub>420</sub>. *M. maripaludis* encodes two copies of *fdhAB* where only *fdh1* is essential for growth on formate and the  $\Delta$ *fdh2* mutant is indistinguishable from wildtype [31<sup>••</sup>,34]. Whole cell transcriptomics and gene specific lacZ fusions have shown that *fdh1* is upregulated during H<sub>2</sub> limitation and in the presence of formate, whereas *fdh2* is only upregulated under formate limiting conditions (Figure 2b) [34]. Likewise, many core methanogenesis genes appear to be regulated in response to formate (Figure 2b) [31<sup>••</sup>]. Whether the regulatory signals for formate and H<sub>2</sub> are the same, somewhat overlapping, or completely distinct is yet to be determined.

### Methanogens with cytochromes

Members of the Order *Methanosarcinales* encode cytochromes and a membrane bound electron transport chain for energy conservation during methanogenesis [1,2,4]. These methanogens are metabolically diverse [4,35<sup>•</sup>], providing a unique opportunity for the characterization of genes involved in the substrate specific regulation of methanogenesis. While many species within the Genus *Methanosarcina* are genetically tractable [36,37], *M. acetivorans* is emerging as a preeminent model system for understanding gene regulation in methanogens and perhaps, even more broadly, in the Archaea. Consequently, *M. acetivorans* encodes the largest number of transcription factors (TFs) in an archaeal genome [14] and substantial work has been done to show that several members of the ArsR family TFs act as both activators and repressors of genes involved in methanogenic metabolism [38,39<sup>••</sup>].

### Methylotrophic methanogenesis

During growth on methylated compounds, also referred to as methylotrophic methanogenesis, 1/4th of the C<sub>1</sub> substrate is oxidized to CO<sub>2</sub> via a reversal of six steps of the core methanogenesis pathway and 3/4th is reduced to methane using the enzyme methyl coenzyme reductase (Figure 3a) [3,40]. During this process, the methyl group from the substrate enters core methanogenesis at the level of methyl-CoM via two stepwise methyl-transfer reactions mediated by two different methyltransferases, referred to as MT1 and MT2 (Figure 3a) [41]. The MT1 complex is substrate-specific and consists of a methyltransferase that transfers the methyl group to its associated corrinoid containing protein, whereas MT2 catalyzes the transfer of the methyl group from the corrinoid protein to CoM to form methyl-CoM (Figure 3a) [42–45]. For the purpose of this review, we


will focus *only* on the regulation of MT1 and MT2 during methylotrophic methanogenesis.

*Methanosarcina* spp. often encode multiple copies of the substrate-specific MT1 that are dynamically regulated (Figure 3b) [18,46]. In *M. acetivorans*, the three copies of methanol-specific MT1 operons, *mtaCB1*, *mtaCB2*, and *mtaCB3* are differentially regulated with respect to changes in growth phase and substrate availability [38,39<sup>••</sup>]. Several members of the ArsR family of TFs that are in the genomic proximity of these operons have been designated as members of the Msr (methanol specific regulator) family upon functional characterization (Figure 3b) [39<sup>••</sup>]. Msr proteins can both activate and repress the expression of *mtaCB* in *M. acetivorans*. Mutational analyses coupled to transcriptional fusions have revealed that MsrA and MsrB act as positive regulators of *mtaCB1* while MsrD and MsrE function in the dual activation and repression of *mtaCB2* and *mtaCB3*, respectively (Figure 3b) [39<sup>••</sup>]. This model of Msr-mediated regulation can be extended to a class of fused MT1/MT2 proteins (MtsD, MtsF, MtsH) that are required for the production of methylated sulfur compounds during growth on carbon monoxide, or for growth on methylated sulfur compounds [47,48]. Genetic studies have shown that MsrF, MsrC, and MsrG are required for the transcriptional activation of MtsD, MtsF, and MtsH respectively (Figure 3b) [49<sup>••</sup>]. While each of the Msr proteins contains a DNA-binding domain, whether these proteins directly bind to the promoter region of the corresponding MTs or interact with other TFs to regulate transcription is unknown and warrants further investigation [39<sup>••</sup>].

Recent work has identified two redox-active cytoplasmic sensor kinases that are likely involved in the regulation of MTs [50,51]. A methylsulfide-specific sensor, named MsmS, is encoded directly upstream of the regulator MsrG and undergoes autophosphorylation when the iron in its covalently bound heme group is oxidized from Fe(II) to Fe(III) (Figure 3b) [50]. Interestingly, a  $\Delta$ *msmS* knockout constitutively produces MtsF, indicating that MsmS may act as a negative effector of *mtsF* expression [50]. Similarly, another sensor kinase, RdmS, is autophosphorylated at a conserved tyrosine residue when intramolecular disulfide bonds between two cysteine residues are formed under oxidizing conditions [51]. These sensor kinases are promising models for understanding archaeal two-component systems, but whether they interact with Msr family regulators to fine-tune the expression of MTs remains unclear.

Beyond transcription initiation, regulation of methylotrophic methanogenesis also occurs at the post-transcriptional level. Curiously, when the *mtaCBA* genes are expressed from a tetracycline-inducible P<sub>*mcrB(tetO1)*</sub> promoter, RNA and protein levels can vary substantially (>10 fold) depending on the growth conditions [52]. This observation is supported by recent work identifying that

Figure 3



Methylo trophic Methanogenesis in *Methanosarcina acetivorans*. **(a)** A substrate-specific methyltransferase (MtaB) catalyzes a methyl-transfer reaction from methanol to a corrinoid-containing protein (MtaC). The heterodimer of MtaB and MtaC (in blue) is called the MT1. A second methyltransferase (MtaA2; in yellow) catalyzes a methyl-transfer reaction from methyl-MtaC to form methyl-coenzyme M (CoM), which is disproportionated in a 3:1 ratio to methane and  $\text{CO}_2$  using the conserved seven-step methanogenesis pathway. The MtaC corrinoid cofactor, methylated 5-hydroxybenzimidazolylcobamide, is depicted in the inset above MT1. **(b)** Msr proteins regulate the expression of three different MtaCB operons in *M. acetivorans*. The dashed arrow indicates that MsrC is a weak activator of MtaCB2 compared to MsrDE.

global changes in mRNA half-life in *M. acetivorans* depend on its growth conditions [53], in addition to work characterizing small RNAs that regulate expression of MTs in *Methanosarcina mazei* [54]. However, a model describing the extent to which post-transcriptional regulation fine tunes gene expression and cellular protein levels has yet to be uncovered.

### Acetoclastic methanogenesis

In *Methanosarcina* spp. growth on acetate proceeds through the dismutation of acetate to CO<sub>2</sub> and methane using two dedicated enzymes: acetate kinase (Ack) and phosphotransacetylase (Pta) [3,55]. First, acetate is activated to acetyl-CoA by Ack and Pta, after which it is split into enzyme-bound methyl and CO groups by acetyl-CoA decarbonylase/synthase (ACDS). The methyl group then enters the central pathway as methyl-tetrahydrosarcinopterin (methyl-H<sub>4</sub>SPT), and the CO group is oxidized to CO<sub>2</sub> [3,55].

Mutational analyses coupled with RNA-seq have revealed that a TrmB family regulator, named MreA, plays a role in the dual activation and repression of acetoclastic and methylotrophic methanogenesis, respectively [56\*\*]. RNA-seq analyses of the  $\Delta mreA$  mutant revealed wholesale downregulation of genes required for acetoclastic methanogenesis with concurrent upregulation of MTs and the Msr family regulatory proteins involved in methylotrophic methanogenesis [56\*\*]. Additionally, purified MreA has been shown to bind the promoter region of *pta* and *fpo* (F<sub>420</sub> dehydrogenase); the former is essential to acetoclastic methanogenesis, while the latter is a membrane bound energy conservation complex upregulated during methylotrophic growth. Taken with the observation of a substantial growth defect in a  $\Delta mreA$  mutant, these results indicate that MreA may play a role in the global regulation of methanogenesis [56\*\*]. However, the underlying mechanism of MreA mediated regulation is yet to be characterized.

### Conclusions and future directions

Methanogenesis is a widespread, environmentally relevant, and evolutionarily important metabolism. While there is considerable phenomenological evidence of regulation in methanogens, a clear picture of transcriptional and post-transcriptional regulatory processes in these pivotal organisms has yet to emerge. The work outlined in this review represents a foundation for future research threads that will shed light on how methanogens sense their environment and tune their metabolism in response. Recent methodological advances, such as genome-wide functional screens coupled with high-throughput global gene expression profiling, are likely to facilitate future efforts to map the complex regulatory network in methanogens. Furthermore, besides its myriad practical applications, studying the regulation of methanogenesis also

presents an opportunity to deepen our fundamental understanding of gene regulation in the domain Archaea.

### Funding information

The authors acknowledge funding from the Searle Scholars Program sponsored by the Kinship Foundation (to D. D.N), the Rose Hills Innovator Grant (to D.D.N), the Beckman Young Investigator Award sponsored by the Arnold and Mabel Beckman Foundation (to D.D.N), the Packard Fellowship in Science and Engineering sponsored by the David and Lucille Packard Foundation (to D.D.N), and startup funds from the Department of Molecular and Cell Biology at UC Berkeley (to D.D.N). The funders had no role in the conceptualization and writing of this review or the decision to submit the work for publication.

### Conflict of interest statement

Nothing declared.

### Acknowledgements

The authors would like to acknowledge members of the Nayak lab for their feedback and input. Molecular graphics were produced using the UCSF Chimera package from the Resources for Biocomputing, Visualization, and Informatics at the University of California, San Francisco (supported by N.I. H.P41 RR-01081).

### References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Thauer RK, Kaster AK, Seedorf H, Buckel W, Hedderich R: **methanogenic archaea: ecologically relevant differences in energy conservation.** *Nat Rev Microbiol* 2008, **6**:579-592.
  2. Borrel G et al.: **Wide diversity of methane and short-chain alkane metabolisms in uncultured archaea.** *Nat Microbiol* 2019, **4**:603-613.
  3. Thauer RK: **Biochemistry of methanogenesis: a tribute to Marjory Stephenson.** *Microbiology* 1998, **144**:2377-2406.
  4. Liu Y, Whitman WB: **Metabolic, phylogenetic, and ecological diversity of the methanogenic archaea.** *Ann N Y Acad Sci* 2008, **1125**:171-189.
  5. Allen G: **Rebalancing the global methane budget.** *Nature* 2016, **538**:46-48.
  6. Reeburgh WS: **Oceanic methane biogeochemistry.** *Chem Rev* 2007, **107**:486-513.
  7. Offre P, Spang A, Schleper C: **Archaea in biogeochemical cycles.** *Annu Rev Microbiol* 2013, **67**:437-457.
  8. Langer D, Hain J, Thuriaux P, Zillig W: **Transcription in archaea: similarity to that in eucarya.** *Proc Natl Acad Sci U S A* 1995, **92**:5768-5772.
  9. Bell SD, Jackson SP: **Mechanism and regulation of transcription in archaea.** *Curr Opin Microbiol* 2001, **4**:208-213.
  10. Hirata A, Klein BJ, Murakami KS: **The X-ray crystal structure of RNA polymerase from archaea.** *Nature* 2008, **451**:851-854.
  11. Kosa PF, Ghosh G, DeDecker BS, Sigler PB: **The 2.1-Å crystal structure of an archaeal preinitiation complex: TATA-box-binding protein/transcription factor (II)B core/TATA-box.** *Proc Natl Acad Sci U S A* 1997, **94**:6042-6047.

## 14 Cell regulation

12. Hausner W, Wettach J, Hethke C, Thomm M: **Two transcription factors related with the eucaryal transcription factors TATA-binding protein and transcription factor IIB direct promoter recognition by an archaeal RNA polymerase.** *J Biol Chem* 1996, **271**:30144-30148.
13. French SL, Santangelo TJ, Beyer AL, Reeve JN: **Transcription and translation are coupled in archaea.** *Mol Biol Evol* 2007, **24**:893-895.
14. Pérez-Rueda E, Janga SC: **Identification and genomic analysis of transcription factors in archaeal genomes exemplifies their functional architecture and evolutionary origin.** *Mol Biol Evol* 2010, **27**:1449-1459.
15. Kim DJ, Forst S: **Genomic analysis of the histidine kinase family in bacteria and archaea.** *Microbiology* 2001, **147**:1197-1212.
16. Galperin MY, Makarova KS, Wolf YI, Koonin EV: **Phyletic distribution and lineage-specific domain architectures of archaeal two-component signal transduction systems.** *J Bacteriol* 2018, **200**:1-16
- Provides an in depth survey of over 2000 histidine kinase and response regulators from 218 archaeal genomes, underlining the differences between archaeal and bacterial two-component systems.
17. Hendrickson EL *et al.*: **Complete genome sequence of the genetically tractable hydrogenotrophic methanogen *Methanococcus maripaludis*.** *J Bacteriol* 2004, **186**:6956-6969.
18. Galagan JE *et al.*: **The genome of *M. acetivorans* reveals extensive metabolic and physiological diversity.** *Genome Res* 2002, **12**:532-542.
19. Imachi H, Sakai S, Nagai H, Yamaguchi T, Takai K: ***Methanofollis ethanolicus* sp. nov., an ethanol-utilizing methanogen isolated from a lotus field.** *Int J Syst Evol Microbiol* 2009, **59**:800-805.
20. Borrel G *et al.*: **Phylogenomic data support a seventh order of methylotrophic methanogens and provide insights into the evolution of methanogenesis.** *Genome Biol Evol* 2013, **5**:1769-1780.
21. Kaster AK, Moll J, Parey K, Thauer RK: **Coupling of ferredoxin and heterodisulfide reduction via electron bifurcation in hydrogenotrophic methanogenic archaea.** *Proc Natl Acad Sci U S A* 2011, **108**:2981-2986.
22. Costa KC *et al.*: **Protein complexing in a methanogen suggests electron bifurcation and electron delivery from formate to heterodisulfide reductase.** *Proc Natl Acad Sci U S A* 2010, **107**:11050-11055.
23. Costa KC, Leigh JA: **Metabolic versatility in methanogens.** *Curr Opin Biotechnol* 2014, **29**:70-75.
24. Fonseca DR, Halim MFA, Holten MP, Costa KC: **Type IV-like pili facilitate transformation in naturally competent archaea.** *J Bacteriol* 2020, **202**:1-12.
25. Susanti D, Frazier MC, Mukhopadhyay B: **A genetic system for *Methanocaldococcus janaschii*: an evolutionarily deeply rooted hyperthermophilic methanoarchaeon.** *Front Microbiol* 2019, **10**:1256.
26. Leigh JA: **Nitrogen fixation in methanogens: the archaeal perspective.** *Curr Issues Mol Biol* 2000, **2**:125-131.
27. Leigh JA, Dodsworth JA: **Nitrogen regulation in bacteria and archaea.** *Annu Rev Microbiol* 2007, **61**:349-377.
28. Karr EA: **Transcription regulation in the third domain.** *Adv Appl Microbiol* 2014, **89**:101-133.
29. Hendrickson EL, Haydock AK, Moore BC, Whitman WB, Leigh JA: **Functionally distinct genes regulated by hydrogen limitation and growth rate in methanogenic archaea.** *Proc Natl Acad Sci U S A* 2007, **104**:8930-8934.
30. Hendrickson EL *et al.*: **Global responses of *Methanococcus maripaludis* to specific nutrient limitations and growth rate.** *J Bacteriol* 2008, **190**:2198-2205.
31. Costa KC *et al.*: **Effects of H<sub>2</sub> and formate on growth yield and regulation of methanogenesis in *Methanococcus maripaludis*.** *J Bacteriol* 2013, **195**:1456-1462
- Identifies genes that are differentially expressed in response to H<sub>2</sub> and formate limitation during continuous culture in defined media.
32. Pihl TD, Sharma S, Reeve JN: **Growth phase-dependent transcription of the genes that encode the two methyl coenzyme M reductase isoenzymes and N<sub>5</sub>-methyltetrahydromethanopterin: coenzyme M methyltransferase in *Methanobacterium thermoautotrophicum* ΔH.** *J Bacteriol* 1994, **176**:6384-6391.
33. Morgan RM, Pihl TD, Nölling J, Reeve JN: **Hydrogen regulation of growth, growth yields, and methane gene transcription in *Methanobacterium thermoautotrophicum* δH.** *J Bacteriol* 1997, **179**:889-898.
34. Wood GE, Haydock AK, Leigh JA: **Function and regulation of the formate dehydrogenase genes of the methanogenic archaeon *Methanococcus maripaludis*.** *J Bacteriol* 2003, **185**:2548-2554.
35. Evans PN *et al.*: **An evolving view of methane metabolism in the archaea.** *Nat Rev Microbiol* 2019, **17**:219-232
- Outlines recent discoveries expanding our understanding of the diversity and distribution of methane metabolism in Archaea with respect to both methanogenesis and anaerobic methanotrophy.
36. Kohler PRA, Metcalf WW: **Genetic manipulation of *Methanosarcina* spp.** *Front Microbiol* 2012, **3**:1-9.
37. Nayak DD, Metcalf WW: **Cas9-mediated genome editing in the methanogenic archaeon *Methanosarcina acetivorans*.** *Proc Natl Acad Sci U S A* 2017, **114**:2976-2981.
38. Bose A, Pritchett MA, Rother M, Metcalf WW: **Differential regulation of the three methanol methyltransferase isozymes in *Methanosarcina acetivorans* C2A.** *J Bacteriol* 2006, **188**:7274-7283.
39. Bose A, Metcalf WW: **Distinct regulators control the expression of methanol methyltransferase isozymes in *Methanosarcina acetivorans* C2A.** *Mol Microbiol* 2008, **67**:649-661
- Characterizes a subset of Msr-family regulators that contribute to the regulation of methanol-specific methyltransferase isozymes, establishing a new class of regulatory proteins in methanogens.
40. Hippe H: **Utilization of trimethylamine and other N-methyl compounds for growth and methane formation by *Methanosarcina barkeri*.** *Proc Natl Acad Sci U S A* 1979, **76**:494-498.
41. Ferry JG: **Enzymology of one-carbon metabolism in methanogenic pathways.** *FEMS Microbiol Rev* 1999, **23**:13-38.
42. van der Meijden P, Te Brommelstroet BW, Poirot CM, van der Drift C, Vogels GD: **Purification and properties of methanol:5-hydroxybenzimidazolylcobamide methyltransferase from *Methanosarcina barkeri*.** *J Bacteriol* 1984, **160**:629-635.
43. Wassenaar RW, Daas PJH, Geerts WJ, Keltjens JT, Van der Drift C: **Involvement of methyltransferase-activating protein and methyltransferase 2 isoenzyme II in methylamine: coenzyme M methyltransferase reactions in *Methanosarcina barkeri* fusaro.** *J Bacteriol* 1996, **178**:6937-6944.
44. Sauer K, Thauer RK: **Methanol: coenzyme M methyltransferase from *Methanosarcina barkeri*. Identification of the active-site histidine in the corrinoid-harboring subunit MtaC by site-directed mutagenesis.** *Eur J Biochem* 1998, **253**:698-705.
45. Ferguson DJ, Krzycki JA, Grahame DA: **Specific roles of methylcobamide: coenzyme M methyltransferase isozymes in metabolism of methanol and methylamines in *Methanosarcina barkeri*.** *J Biol Chem* 1996, **271**:5189-5194.
46. Maeder DL *et al.*: **The *Methanosarcina barkeri* genome: comparative analysis with *Methanosarcina acetivorans* and *Methanosarcina mazei* reveals extensive rearrangement within *Methanosarcinal* genomes.** *J Bacteriol* 2006, **188**:7922-7931.
47. Oelgeschläger E, Rother M: **In vivo role of three fused corrinoid/methyl transfer proteins in *Methanosarcina acetivorans*.** *Mol Microbiol* 2009, **72**:1260-1272.
48. Lessner DJ *et al.*: **An unconventional pathway for reduction of CO<sub>2</sub> to methane in CO-grown *Methanosarcina acetivorans***

- revealed by proteomics. *Proc Natl Acad Sci U S A* 2006, **103**:17921-17926.
49. Bose A, Kulkarni G, Metcalf WW: **Regulation of putative methylsulphide methyltransferases in *Methanosarcina acetivorans* C2A.** *Mol Microbiol* 2009, **74**:227-238  
 Identifies and characterizes novel Msr-family regulators that influence expression of methylsulfide methyltransferases, expanding the role of this class of archaea-specific regulators.
  50. Molitor B *et al.*: **A heme-based redox sensor in the methanogenic archaeon *Methanosarcina acetivorans*.** *J Biol Chem* 2013, **288**:18458-18472.
  51. Fiege K, Frankenberg-Dinkel N: **Thiol-based redox sensing in the methyltransferase associated sensor kinase RdmS in *Methanosarcina acetivorans*.** *Environ Microbiol* 2019, **21**:1597-1610.
  52. Oplencia RB, Bose A, Metcalf WW: **Physiology and posttranscriptional regulation of methanol: coenzyme M methyltransferase isozymes in *Methanosarcina acetivorans* C2A.** *J Bacteriol* 2009, **191**:6928-6935.
  53. Peterson JR *et al.*: **Genome-wide gene expression and RNA half-life measurements allow predictions of regulation and metabolic behavior in *Methanosarcina acetivorans*.** *BMC Genomics* 2016, **17**:1-23.
  54. Jäger D, Pernitzsch SR, Richter AS, Backofen R, Sharma CM, Schmitz RA: **An archaeal sRNA targeting *cis*- and *trans*-encoded mRNAs via two distinct domains.** *Nucleic Acids Res* 2012, **40**:10964-10979.
  55. Jetten MSM, Stams AJM, Zehnder AJB: **Methanogenesis from acetate: a comparison of the acetate metabolism in *Methanotherix soehngenii* and *Methanosarcina* spp.** *FEMS Microbiol Lett* 1992, **88**:181-197.
  56. Reichlen MJ, Vepachedu VR, Murakami KS, Ferry JG: **MreA functions in the global regulation of methanogenic pathways in *Methanosarcina acetivorans*.** *mBio* 2012, **3**:1-9  
 Two transcription factors identifies MreA, a TrmB family regulator, that plays a role in the activation of acetoclastic methanogenesis and repression of methylotrophic methanogenesis.